

# the MSR-INRIA Joint Centre

*Jean-Jacques Lévy*

*IKI, June 11, 2008*

CENTRE DE RECHERCHE  
COMMUN



# Plan


## 1. Context

## 2. Track A

- Math. Components
- Security
- TLA+

## 3. Track B

- DDMF
- ReActivity
- Adaptative search
- Image & video mining



# Context

# Management

INRIA



Gilles Kahn

Michel Cosnard

Michel Bidoit  
Bruno Sportisse



MSR Cambridge



Roger Needham

Andrew Herbert

Bernard Ourghanian  
Thomas Serval


Eric Boussouler  
Stephen Emmott  
Gérard Giraudon  
Gérard Huet  
Marc Jalabert  
Jean Vuillemin  
Ken Wood

Joint  
Centre

J.-J. Lévy

Andrew Blake  
Stephen Emmott  
Malik Ghallab  
Claude Puech

# Long cooperation among researchers



# Organization


## a rather complex system

- **7 research projects (in two tracks)**
- **12 resident researchers**
- **non permanent researchers funded by the Joint Centre**
- **permanent researchers paid by INRIA or MSR**
- **operational support by INRIA Saclay**
- **1 system manager** (Guillaume Rousse, INRIA Saclay)
- **1 administrative assistant** (Martine Thirion, Joint Centre)
- **1 deputy director** (Pierre-Louis Xech, MS France)
- **active support from MS France**

# Localization

## the plateau de Saclay

= long term  
investment





# People

## PhD Students

- Francois GARILLOT
- Sidi OULD BIHA
- Iona PASCA
- Roland ZUMKELLER
- Pierre-Malo DENIELOU
- Nataliya GUTS
- Jérémie PLANUL
- Santiago ZANELLA
- Alexandre BENOIT
- Marc MEZZAROBA
- Nathalie HENRY (+)
- Nicolas MASSON
- Arnaud SPIVAK
- Aurélien TABARD

## Post Docs

- Alejandro ARBALAEZ
- Alvaro FIALHO
- Adrien GAIDON
- Stéphane LE ROUX
- Guillaume MELQUIOND (\*)
- Assia MAHBOUBI (\*)
- Ricardo CORIN (\*)
- Gurvan LE GUERNIC
- Eugen ZALINESCU
- Tamara REZK (\*)
- Kaustuv CHAUDURI (\*?)
- Stefan GERHOLD
- Fanny CHEVALIER
- Niklas ELMQVIST
- Catherine LEDONTAL
- Tomer MOSCOVICH
- Theophanis TSANDILAS
- Nikolaus HANSEN (\*?)
- Neva CHERNIAVSKY

---

(\*) Now on permanent INRIA position, (+) on permanent MSR position

# Track A

*Software Security  
Trustworthy Computing*

# Mathematical components

Georges Gonthier, MSRC

Assia Mahboubi, INRIA Saclay/LIX

Andrea Asperti, Bologna

Y. Bertot, L. Rideau, L. Théry, Sidi Ould Biha,

Iona Pasca, INRIA Sophia

François Garillot, MSR-INRIA (PhD)

Guillaume Melquiond, MSR-INRIA (postdoc)


Stéphane le Roux, MSR-INRIA (postdoc)

Benjamin Werner, INRIA Saclay/LIX,

Roland Zumkeller, LIX (PhD)

## Computational proofs

- computer assistance for long formal proofs.
- reflection of computations into Coq-logic: [ssreflect](#).



4-color

Appel-Haken



finite groups

Feit-Thompson



Kepler

Hales

## Section R\_props.

(\* The **ring** axioms, and some useful basic corollaries. \*)

```
Hypothesis mult1x : forall x, 1 * x = x.
Hypothesis mult0x : forall x : R, 0 * x = 0.
Hypothesis plus0x : forall x : R, 0 + x = x.
Hypothesis minusxx : forall x : R, x - x = 0.
Hypothesis plusA : forall x1 x2 x3 : R, x1 + (x2 + x3) = x1 + x2 + x3.
Hypothesis plusC : forall x1 x2 : R, x1 + x2 = x2 + x1.
Hypothesis multA : forall x1 x2 x3 : R, x1 * (x2 * x3) = x1 * x2 * x3.
Hypothesis multC : forall x1 x2 : R, x1 * x2 = x2 * x1.
Hypothesis distrR : forall x1 x2 x3 : R, (x1 + x2) * x3 = x1 * x3 + x2 * x3.
```

Lemma plusCA : forall x1 x2 x3 : R, x1 + (x2 + x3) = x2 + (x1 + x3).

Proof. move=> \*; rewrite !plusA; congr (\_ + \_); exact: plusC. Qed.

Lemma multCA : forall x1 x2 x3 : R, x1 \* (x2 \* x3) = x2 \* (x1 \* x3).

Proof. move=> \*; rewrite !multA; congr (\_ \* \_); exact: multC. Qed.

Lemma distrL : forall x1 x2 x3 : R, x1 \* (x2 + x3) = x1 \* x2 + x1 \* x3.

Proof. by move=> x1 x2 x3; rewrite !(multC x1) distrR. Qed.

Lemma oppK : involutive opp.

Proof.

by move=> x; rewrite -{2}[x]plus0x -(minusxx (- x)) plusC plusA minusxx plus0x.

Qed.

Lemma multm1x : forall x, -1 \* x = -x.

Proof.

move=> x; rewrite -[\_ \* x]plus0x -(minusxx x) -{1}[x]mult1x plusC plusCA plusA.

by rewrite -distrR minusxx mult0x plus0x.

Qed.

Lemma Rsubn : forall m n, m >= n -> (m - n)%N = m - n :> R.  
 Proof.  
~~move=> m n; move/leq\_add\_sub=> Dm.~~  
 by rewrite -{2}Dm Raddn -plusA plusCA minusxx plusC plus0x.  
 Qed.

Lemma Rmuln : forall m n, (m \* n)%N = m \* n :> R.  
 Proof.  
~~move=> m n; elim: m => /= [!m IHm]; first by rewrite mult0x.~~  
 by rewrite Raddn RofSnE IHm distrR mult1x plusC.  
 Qed.

Lemma RexpSnE : forall x n, RexpSn x n = x ^ n \* x.  
 Proof. by move=> x; elim=> /= [!\_ -> //]; rewrite mult1x. Qed.

Lemma mult\_exp : forall x1 x2 n, (x1 \* x2) ^ n = x1 ^ n \* x2 ^ n.  
 Proof.  
~~by move=> x1 x2; elim=> //=\_IHn; rewrite !RexpSnE IHn -!multA (multCA x1).~~  
 Qed.

Lemma exp\_addn : forall x n1 n2, x ^ (n1 + n2) = x ^ n1 \* x ^ n2.  
 Proof.  
~~move=> x n1 n2; elim: n1 => /= [!n1 IHn]; first by rewrite mult1x.~~  
 by rewrite !RexpSnE IHn multC multCA multA.  
 Qed.

Lemma Rexpn : forall m n, (m ^ n)%N = m ^ n :> R.  
 Proof. by move=> m; elim=> //=\_IHn; rewrite Rmuln RexpSnE IHn multC. Qed.

Lemma exp0n : forall n, 0 < n -> 0 ^ n = 0.  
 Proof. by move=> [\_[\_n]] //=\_; rewrite multC mult0x. Qed.

Lemma exp1n : forall n, 1 ^ n = 1.  
 Proof. by elim=> //=\_IHn; rewrite RexpSnE IHn mult1x. Qed.

Lemma exp\_mulin : forall x n1 n2, x ^ (n1 \* n2) = (x ^ n1) ^ n2.  
 Proof.  
~~move=> x n1 n2; rewrite mulnC; elim: n2 => //=\_n2 IHn.~~  
 by rewrite !RexpSnE exp\_addn IHn multC.  
 Qed.

Lemma sign\_odd : forall n, (-1) ^ odd n = (-1) ^ n.  
 Proof.  
~~move=> n; rewrite -{2}[n]odd\_double\_half addnC double\_mul2 exp\_addn exp\_mulin.~~  
 by rewrite /= mult1x oppK expin mult1x.  
 Qed.

Lemma sign\_adbb : forall b1 b2, (-1) ^ (b1 (+) b2) = (-1) ^ b1 \* (-1) ^ b2.  
 Proof. by do 2!case; rewrite //=\_?mult1x ?mult1x ?oppK. Qed. □

Lemma sign\_permM : forall d (s t : permType d),  
 -(DOS)-- determinant.v 45% (760,61) (coq)

rewrite isum0 ?plus0x // => i'; rewrite andbT; move/negbET->; exact: mult0x.  
Qed.

Lemma matrix\_transpose\_mul : forall m n p (A : M\_(m, n)) (B : M\_(n, p)),  
  \At (A \*m B) =m \At B \*m \At A.  
Proof. split=> k i; apply: eq\_isumR => j \_; exact: multC. Qed.

Lemma matrix\_multx1 : forall m n (A : M\_(m, n)), A \*m \1m =m A.  
Proof.

move=> m n A; apply: matrix\_transpose\_inj.  
by rewrite matrix\_transpose\_mul matrix\_transpose\_unit matrix\_mult1x.  
Qed.

Lemma matrix\_distrR : forall m n p (A1 A2 : M\_(m, n)) (B : M\_(n, p)),  
  (A1 +m A2) \*m B =m A1 \*m B +m A2 \*m B.

Proof.  
move=> m n p A1 A2 B; split=> i k /; rewrite -isum\_plus.  
by apply: eq\_isumR => j \_; rewrite -distrR.  
Qed.

Lemma matrix\_distrL : forall m n p (A : M\_(m, n)) (B1 B2 : M\_(n, p)),  
  A \*m (B1 +m B2) =m A \*m B1 +m A \*m B2.

Proof.  
move=> m n p A B1 B2; apply: matrix\_transpose\_inj.  
rewrite matrix\_transpose\_plus !matrix\_transpose\_mul.  
by rewrite -matrix\_distrR -matrix\_transpose\_plus.  
Qed.

Lemma matrix\_multA : forall m n p q  
  (A : M\_(m, n)) (B : M\_(n, p)) (C : M\_(p, q)), □  
  A \*m (B \*m C) =m A \*m B \*m C.

Proof.  
move=> m n p q A B C; split=> i l /=.  
transitivity (\sum\_(k) (\sum\_(j) (A i j \* B j k \* C k l))).  
  rewrite exchange\_isum; apply: eq\_isumR => j \_; rewrite isum\_distrL.  
  by apply: eq\_isumR => k \_; rewrite multA.  
by apply: eq\_isumR => j \_; rewrite isum\_distrR.  
Qed.

Lemma perm\_matrixM : forall n (s t : S\_(n)),  
  perm\_matrix (s \* t)%G =m perm\_matrix s \*m perm\_matrix t.

Proof.  
move=> n; split=> i j /=; rewrite (isumD1 (s i)) // set11 mult1x -permM.  
rewrite isum0 => [lj']; first by rewrite plusC plus0x.  
by rewrite andbT; move/negbET->; rewrite mult0x.  
Qed.

Lemma matrix\_trace\_plus : forall n (A B : M\_(n)), \tr (A +m B) = \tr A + \tr B.  
Proof. by move=> n A B; rewrite -isum\_plus. Qed.

Lemma matrix\_trace\_scale : forall n x (A : M\_(n)), \tr (x \*sm A) = x \* \tr A.  
Proof. by move=> \*; rewrite isum\_distrL. Qed.

-(DOS)-- determinant.v 77% (1190,48) (coq)

(\* And now, finally, the title feature. \*)

```
Lemma determinant_multilinear : forall n (A B C : M_(n)) i0 b c,
  row i0 A =m b *sm row i0 B +m c *sm row i0 C ->
  row' i0 B =m row' i0 A -> row' i0 C =m row' i0 A ->
  \det A = b * \det B + c * \det C.
```

Proof.

move=> n A B C i0 b c ABC.

move/matrix\_eq\_rem\_row=> BA; move/matrix\_eq\_rem\_row=> CA.

rewrite !isum\_distrL -isum\_plus; apply: eq\_isumR => s ..

rewrite !(multCA (\_ ^ s)) -distrL; congr (\_ \* \_).

rewrite !(@iprodD1 \_ i0 (setA \_)) // (matrix\_eq\_row ABC) distrR !multA.

by congr (\_ \* \_ + \_ \* \_); apply: eq\_iprodR => i;

rewrite andbT => ?; rewrite ?BA ?CA.

Qed.

```
Lemma alternate_determinant : forall n (A : M_(n)) i1 i2,
  i1 != i2 -> A i1 =1 A i2 -> \det A = 0.
```

Proof.

move=> n A i1 i2 Di12 A12; pose r := I\_(n).

pose t := transp i1 i2; pose tr s := (t \* s)%G.

have trk : involutive tr by move=> s; rewrite /tr multgA transp2 mulig.

have Etr: forall s, odd\_perm (tr s) = even\_perm s.

by move=> s; rewrite odd\_permM odd\_transp Di12.

rewrite /(\det \_) (isumID (@even\_perm r)) /%; set S1 := \sum\_(in \_) ..

rewrite -{2}(minusxx S1); congr (\_ + \_); rewrite {}//S1 -isum\_opp.

rewrite (reindex\_isum tr); last by exists tr.

symmetry; apply: eq\_isum => [s | s seven]; first by rewrite negbK Etr.

rewrite -multmix multA Etr seven (negbET seven) multmix; congr (\_ \* \_).

rewrite (reindex\_iprod t); last by exists (t : \_ -> \_) => i \_; exact: transpK.

apply: eq\_iprodR => i \_; rewrite permM /t.

by case: transpP => // ->; rewrite A12.

Qed. □

```
Lemma determinant_transpose : forall n (A : M_(n)), \det (\^t A) = \det A.
```

Proof.

move=> n A; pose r := I\_(n); pose ip p : permType r := p^-1.

rewrite /(\det \_) (reindex\_isum ip) /%; last first.

by exists ip => s \_; rewrite /ip invgK.

apply: eq\_isumR => s \_; rewrite odd\_permV / (reindex\_iprod s).

by congr (\_ \* \_); apply: eq\_iprodR => i \_; rewrite permK.

by exists (s^-1 : \_ -> \_) => i \_; rewrite ?permK ?permKv.

Qed.

```
Lemma determinant_perm : forall n s, \det (@perm_matrix n s) = (-1) ^ s.
```

Proof.

move=> n s; rewrite /(\det \_) (isumD1 s) //.

rewrite iprod1 => [!i \_]; last by rewrite /= set11.

rewrite isum0 => [!t Dst]; first by rewrite plusC plus0x multC mult1x.

case: (pickP (fun i => s i != t i)) => [i ist | Est].

by rewrite (iprodD1 i) // multCA /= (negbET ist) mult0x.

move: Dst; rewrite andbT; case/eqP.

-(DOS)-- determinant.v 81% (1256,4) (coq)

```
Lemma determinant1 : forall n, \det (unit_matrix n) = 1.
```

Proof.

```
move=> n; have:= @determinant_perm n 1%G; rewrite odd_perm1 => /= <-.
```

```
apply: determinant_extensional; symmetry; exact: perm_matrix1.
```

Qed.

□

```
Lemma determinant_scale : forall n x (A : M_(n)),
```

```
\det (x *sm A) = x ^ n * \det A.
```

Proof.

```
move=> n x A; rewrite isum_distrL; apply: eq_isumR => s _.
```

```
by rewrite multCA iprod_mult iprod_id card_ordinal.
```

Qed.

```
Lemma determinantM : forall n (A B : M_(n)), \det (A *m B) = \det A * \det B.
```

Proof.

```
move=> n A B; rewrite isum_distrR.
```

```
pose AB (f : F_(n)) (s : S_(n)) i := A i (f i) * B (f i) (s i).
```

```
transitivity (\sum_(f) \sum_(s : S_(n)) (-1) ^ s * \prod_(i) AB f s i).
```

```
rewrite exchange_isum; apply: eq_isumR => s _.
```

```
by rewrite -isum_distrL distr_iprodA_isumA.
```

```
rewrite (isumID (fun f => uniq (fval f))) plusC isum0 ?plus0x => /= [if Uf].
```

```
rewrite (reindex_isum (fun s => val (pval s))); last first.
```

```
have s0 : S_(n) := 1%G; pose uf (f : F_(n)) := uniq (fval f).
```

```
pose pf f := if insub uf f is Some s then Perm s else s0.
```

```
exists pf => /= f Uf; rewrite /pf (insubT uf Uf) //; exact: eq_fun_of_perm.
```

```
apply: eq_isum => [s|s _]; rewrite ?(valP (pval s)) // isum_distrL.
```

```
rewrite (reindex_isum (mulg s)); last first.
```

```
by exists (mulg s^-1) => t; rewrite ?mulKgv ?mulKg.
```

# Secure Distributed Computations and their Proofs

Cédric Fournet, MSRC

Karthik Bhargavan, MSRC

Ricardo Corin, INRIA Rocq.

Pierre-Malo Deniélo, INRIA Rocq.

G. Barthe, B. Grégoire, S. Zanella, INRIA Sophia

James Leifer, INRIA Rocq.

Jean-Jacques Lévy, INRIA Rocq.

Tamara Rezk, INRIA Sophia

Francesco Zappa Nardelli, INRIA Rocq.

Nataliya Guts, MSR-INRIA (PhD)

Jérémie Planul, MSR-INRIA (intern)

## Distributed computations + Security

- programming with secured communications
- certified compiler from high-level primitives to low-level crypto-protocols
- formal proofs of probabilistic protocols



# Secure Distributed Computations and their Proofs

## Recent results:

- **secure sessions** v2 (proofs by typing)
- concurrent secure sessions v1
- correctness proofs of TLS implementations
- information flow + cryptography
- secure logs
- secure modeling of e-cash

# Tools for formal proofs

Damien Doligez, INRIA Rocq.


Kaustuv Chaudhury, MSR-INRIA (postdoc)

Leslie Lamport, MSRSV


Stephan Merz, INRIA Lorraine

## Natural proofs

- first-order set theory + temporal logic
- specification/verification of concurrent programs.
- tools for automatic theorem proving



TLA+



tools for proofs



Zenon

EXTENDS Naturals

---

(\* First some general logical axioms pulled from the trusted base \*)

(\* The following is a specific instance of a theorem provable by the Peano axioms \*)

THEOREM TwoIsNotOne =

2 # 1

PROOF OMITTED

THEOREM NegElim =

ASSUME

NEW CONSTANT A,

A, ~A

PROVE

FALSE

PROOF OMITTED

THEOREM ImplIntro =

ASSUME

NEW CONSTANT A, NEW CONSTANT B,

ASSUME A PROVE B

PROVE A  $\Rightarrow$  B

PROOF OMITTED

```
(**  
 * Main theorem: there is no irreducible rational number x/y whose  
 * square is 2.  
 *)
```

```
THEOREM SqrtTwoIrrational =
```

```
 \A x, y \in Nat : Coprime(x, y)  $\Rightarrow$  x^2  $\neq$  2 * y^2
```

```
PROOF <1>1. ASSUME
```

```
 NEW x \in Nat,
```

```
 NEW y \in Nat,
```

```
 coprimality::: Coprime(x, y),
```

```
 main::: x^2 = 2 * y^2
```

```
PROVE
```

```
 FALSE
```

```
PROOF <2>1. Divides(2, x)
```

```
 PROOF <3>1. Divides(2, x^2)
```

```
 BY <1>1!3
```

```
<3>2. QED
```

```
 BY <3>1, TwoIsPrime, SquareLemma
```

```
<2>2. Divides(2, y)
```

```
 PROOF <3>1. PICK r \in Nat : x = 2 * r
```

```
 BY <2>1, DivLemma
```

```
<3>2. x^2 = 2 * (2 * r^2)
```

```
 BY <3>1
```

```
<3>3. 2 * y^2 = 2 * (2 * r^2)
```

```
 BY <1>1!main, <3>2
```

```
<3>4. y^2 = 2 * r^2
```

```

<3>2. QED
      BY <3>1, TwoIsPrime, SquareLemma
<2>2. Divides(2, y)
      PROOF <3>1. PICK r \in Nat : x = 2 * r
          BY <2>1, DivLemma
          <3>2. x^2 = 2 * (2 * r^2)
          BY <3>1
          <3>3. 2 * y^2 = 2 * (2 * r^2)
          BY <1>1!main, <3>2
          <3>4. y^2 = 2 * r^2
          BY <3>2, LeftCancellationLemma
          <3>5. QED
              BY <3>3, TwoIsPrime, SquareLemma
<2>3. ~ (Divides(2, y))
      PROOF <3>1. \A d \in Nat : (Divides(d, x) \wedge Divides(d, y)) \Rightarrow d = 1
          BY <1>1!coprimality
          <3>2. 2 = 1
          BY <2>1, <2>2, <3>1
          <3>3. QED
              BY <3>2, TwoIsNotOne
          <3>4. QED
              BY <2>2, <2>3, NegElim
<1>2. QED
      BY <1>1, ImplIntro, ForallIntro

```

(\* The main definitions and lemmas (proofs omitted) \*)

Divides(d, n) =  
   $\wedge d \in \text{Nat}$   
   $\wedge n \in \text{Nat}$   
   $\wedge \exists q \in \text{Nat} : n = d * q$

THEOREM DivLemma =  
   $\forall d, n \in \text{Nat} : \text{Divides}(d, n) \Rightarrow \exists r \in \text{Nat} : n = r * d$

PROOF OMITTED

Prime(x) =  
   $\wedge x \in \text{Nat}$   
   $\wedge \forall d \in \text{Nat} : \text{Divides}(d, x) \Rightarrow \begin{cases} \vee d = 1 \\ \vee d = x \end{cases}$

PrimeNat = {x ∈ Nat : Prime(x)}

THEOREM TwoIsPrime = 2 ∈ PrimeNat

PROOF OMITTED

THEOREM SquareLemma =  
   $\forall p \in \text{PrimeNat}, x \in \text{Nat} :$   
     $\text{Divides}(p, x^2) \Rightarrow \text{Divides}(p, x)$

PROOF OMITTED

# Tools for formal proofs

## Recent results:

- **Proof Manager** with incremental, non-linear proofs
- declarative meta-language
- proofs like done by Mathematicians
- proof of the atomic Bakery algorithm with PM

# Track B

*Computational Sciences  
Scientific Information Interaction*

CENTRE DE RECHERCHE  
COMMUN



# Dynamic dictionary of math functions


Bruno Salvy, INRIA Rocq.,  
Alin Bostan, INRIA Rocq.,  
Frédéric Chyzak, INRIA Rocq.

Henry Cohn, [Theory Group] MSRR  
Alexandre Benoit, MSR-INRIA (intern)  
Marc Mezzarobba, MSR-INRIA (intern)

## Computer Algebra and Web for useful functions,

- dynamic tables of their properties.
- generation of programs to compute them.

**Maple™ 11**



CENTRE DE RECHERCHE  
COMMUN

INRIA  
MICROSOFT RESEARCH

## 9. Bessel Functions of Integer Order

### Mathematical Properties

#### Notation

The tables in this chapter are for Bessel functions of integer order; the text treats general orders. The conventions used are:

$$z = x + iy; \quad x, y \text{ real.}$$

$n$  is positive integer or zero.

$\nu, \mu$  are unrestricted except where otherwise indicated;  $\nu$  is supposed real in the sections devoted to Kelvin functions 9.9, 9.10, and 9.11.

The notation used for the Bessel functions is that of Watson [9.15] and the British Association and Royal Society Mathematical Tables. The function  $Y_\nu(z)$  is often denoted  $N_\nu(z)$  by physicists and European workers.

Other notations are those of:

Aldis, Airey:

$$G_n(z) \text{ for } -\frac{1}{2}\pi Y_n(z), K_n(z) \text{ for } (-)^n K_n(z).$$

Clifford:

$$C_n(x) \text{ for } x^{-\frac{1}{2}n} J_n(2\sqrt{x}).$$

Gray, Mathews and MacRobert [9.9]:

$$Y_n(z) \text{ for } \frac{1}{2}\pi Y_n(z) + (\ln 2 - \gamma) J_n(z),$$

$$\bar{Y}_n(z) \text{ for } \pi e^{\nu\pi i} \sec(\nu\pi) Y_n(z),$$

$$G_\nu(z) \text{ for } \frac{1}{2}\pi H_\nu^{(1)}(z).$$

Jahnke, Emde and Lösch [9.32]:

$$\Lambda_\nu(z) \text{ for } \Gamma(\nu+1)(\frac{1}{2}z)^{-\nu} J_\nu(z).$$

Jeffreys:

$$H_{\nu s}(z) \text{ for } H_\nu^{(1)}(z), H_{\nu i}(z) \text{ for } H_\nu^{(2)}(z),$$

$$Kh_\nu(z) \text{ for } (2/\pi)K_\nu(z).$$

Heine:

$$K_n(z) \text{ for } -\frac{1}{2}\pi Y_n(z).$$

Neumann:

$$Y^n(z) \text{ for } \frac{1}{2}\pi Y_n(z) + (\ln 2 - \gamma) J_n(z).$$

Whittaker and Watson [9.18]:

$$K_\nu(z) \text{ for } \cos(\nu\pi) K_\nu(z).$$

### Bessel Functions $J$ and $Y$

#### 9.1. Definitions and Elementary Properties

##### Differential Equation

$$9.1.1 \quad z^2 \frac{d^2 w}{dz^2} + z \frac{dw}{dz} + (z^2 - \nu^2) w = 0$$

Solutions are the Bessel functions of the first kind  $J_\pm(z)$ , of the second kind  $Y_\nu(z)$  (also called Weber's function) and of the third kind  $H_\nu^{(1)}(z), H_\nu^{(2)}(z)$  (also called the Hankel functions). Each is a regular (holomorphic) function of  $z$  throughout the  $z$ -plane cut along the negative real axis, and for fixed  $z(z \neq 0)$  each is an entire (integral) function of  $\nu$ . When  $\nu = \pm n$ ,  $J_\nu(z)$  has no branch point and is an entire (integral) function of  $z$ .

Important features of the various solutions are as follows:  $J_\nu(z)$  ( $\Re \nu \geq 0$ ) is bounded as  $z \rightarrow 0$  in any bounded range of  $\arg z$ .  $J_\nu(z)$  and  $J_{-\nu}(z)$  are linearly independent except when  $\nu$  is an integer.  $J_\nu(z)$  and  $Y_\nu(z)$  are linearly independent for all values of  $\nu$ .

$H_\nu^{(1)}(z)$  tends to zero as  $|z| \rightarrow \infty$  in the sector  $0 < \arg z < \pi$ ;  $H_\nu^{(2)}(z)$  tends to zero as  $|z| \rightarrow \infty$  in the sector  $-\pi < \arg z < 0$ . For all values of  $\nu$ ,  $H_\nu^{(1)}(z)$  and  $H_\nu^{(2)}(z)$  are linearly independent.

##### Relations Between Solutions

$$9.1.2 \quad Y_\nu(z) = \frac{J_\nu(z) \cos(\nu\pi) - J_{-\nu}(z)}{\sin(\nu\pi)}$$

The right of this equation is replaced by its limiting value if  $\nu$  is an integer or zero.

##### 9.1.3

$$H_\nu^{(1)}(z) = J_\nu(z) + iY_\nu(z) \\ = i \csc(\nu\pi) \{ e^{-\nu\pi i} J_\nu(z) - J_{-\nu}(z) \}$$

##### 9.1.4

$$H_\nu^{(2)}(z) = J_\nu(z) - iY_\nu(z) \\ = i \csc(\nu\pi) \{ J_{-\nu}(z) - e^{\nu\pi i} J_\nu(z) \}$$

$$9.1.5 \quad J_{-n}(z) = (-)^n J_n(z) \quad Y_{-n}(z) = (-)^n Y_n(z)$$

$$9.1.6 \quad H_{-\nu}^{(1)}(z) = e^{\nu\pi i} H_\nu^{(1)}(z) \quad H_{-\nu}^{(2)}(z) = e^{-\nu\pi i} H_\nu^{(2)}(z)$$




FIGURE 9.1.  $J_0(x), Y_0(x), J_1(x), Y_1(x)$ .




FIGURE 9.2.  $J_{10}(x), Y_{10}(x)$ , and  $M_{10}(x) = \sqrt{J_{10}^2(x) + Y_{10}^2(x)}$ .




FIGURE 9.4. Contour lines of the modulus and phase of the Hankel Function  $H_0^{(1)}(x+iy) = M_0 e^{i\phi_0}$ . From E. Jahnke, F. Emde, and F. Lösch, Tables of higher functions, McGraw-Hill Book Co., Inc., New York, N.Y., 1960 (with permission).

## Limiting Forms for Small Arguments

When  $\nu$  is fixed and  $z \rightarrow 0$

9.1.7

$$J_\nu(z) \sim (\frac{1}{2}z)^\nu / \Gamma(\nu+1) \quad (\nu \neq -1, -2, -3, \dots)$$

$$9.1.8 \quad Y_0(z) \sim -iH_0^{(1)}(z) \sim iH_0^{(2)}(z) \sim (2/\pi) \ln z$$

9.1.9

$$Y_\nu(z) \sim -iH_\nu^{(1)}(z) \sim iH_\nu^{(2)}(z) \sim -(1/\pi) \Gamma(\nu) (\frac{1}{2}z)^{-\nu} \quad (\Re \nu > 0)$$

## Ascending Series

$$9.1.10 \quad J_\nu(z) = (\frac{1}{2}z)^\nu \sum_{k=0}^{\infty} \frac{(-\frac{1}{4}z^2)^k}{k! \Gamma(\nu+k+1)}$$

9.1.11

$$Y_n(z) = -\frac{(\frac{1}{2}z)^{-n}}{\pi} \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} (\frac{1}{4}z^2)^k + \frac{2}{\pi} \ln(\frac{1}{2}z) J_n(z) - \frac{(\frac{1}{2}z)^n}{\pi} \sum_{k=0}^{\infty} \{ \psi(k+1) + \psi(n+k+1) \} \frac{(-\frac{1}{4}z^2)^k}{k!(n+k)!}$$

where  $\psi(n)$  is given by 6.3.2.

$$9.1.12 \quad J_0(z) = 1 - \frac{\frac{1}{2}z^2}{(1!)^2} + \frac{(\frac{1}{2}z^2)^2}{(2!)^2} - \frac{(\frac{1}{2}z^2)^3}{(3!)^2} + \dots$$

9.1.13

$$Y_0(z) = \frac{2}{\pi} \{ \ln(\frac{1}{2}z) + \gamma \} J_0(z) + \frac{2}{\pi} \{ \frac{\frac{1}{2}z^2}{(1!)^2} - (1+\frac{1}{2}) \frac{(\frac{1}{2}z^2)^2}{(2!)^2} + (1+\frac{1}{2}+\frac{1}{3}) \frac{(\frac{1}{2}z^2)^3}{(3!)^2} - \dots \}$$

9.1.14

$$J_\nu(z) J_\mu(z) =$$

$$\sum_{k=0}^{\infty} \frac{(-)^k \Gamma(\nu+\mu+2k+1) (\frac{1}{4}z^2)^k}{\Gamma(\nu+k+1) \Gamma(\mu+k+1) \Gamma(\nu+\mu+k+1) k!}$$

## Wronskians

$$W\{J_\nu(z), J_{-\nu}(z)\} = J_{\nu+1}(z) J_{-\nu}(z) + J_\nu(z) J_{-(\nu+1)}(z) = -2 \sin(\nu\pi)/(\pi z)$$

9.1.16

$$W\{J_\nu(z), Y_\nu(z)\} = J_{\nu+1}(z) Y_\nu(z) - J_\nu(z) Y_{\nu+1}(z) = 2/(\pi z)$$

9.1.17

$$W\{H_\nu^{(1)}(z), H_\nu^{(2)}(z)\} = H_{\nu+1}^{(1)}(z) H_\nu^{(2)}(z) - H_\nu^{(1)}(z) H_{\nu+1}^{(2)}(z) = -4i/(\pi z)$$

## Integral Representations

$$9.1.18 \quad J_0(z) = \frac{1}{\pi} \int_0^\pi \cos(z \sin \theta) d\theta = \frac{1}{\pi} \int_0^\pi \cos(z \cos \theta) d\theta$$

9.1.19

$$Y_0(z) = \frac{4}{\pi^2} \int_0^{\frac{\pi}{2}} \cos(z \cos \theta) \{ \gamma + \ln(2z \sin^2 \theta) \} d\theta$$

9.1.20

$$J_\nu(z) = \frac{(\frac{1}{2}z)^\nu}{\pi^4 \Gamma(\nu + \frac{3}{2})} \int_0^\pi \cos(z \cos \theta) \sin^{2\nu} \theta d\theta = \frac{2(\frac{1}{2}z)^\nu}{\pi^4 \Gamma(\nu + \frac{3}{2})} \int_0^1 (1-t^2)^{\nu-\frac{1}{2}} \cos(zt) dt \quad (\Re \nu > -\frac{1}{2})$$

9.1.21

$$J_n(z) = \frac{1}{\pi} \int_0^\pi \cos(z \sin \theta - n\theta) d\theta = \frac{i^{-n}}{\pi} \int_0^\pi e^{iz \cos \theta} \cos(n\theta) d\theta$$

9.1.22

$$J_\nu(z) = \frac{1}{\pi} \int_0^\pi \cos(z \sin \theta - \nu\theta) d\theta - \frac{\sin(\nu\pi)}{\pi} \int_0^\infty e^{-z \sinh t - \nu t} dt \quad (|\arg z| < \frac{1}{2}\pi)$$

$$Y_\nu(z) = \frac{1}{\pi} \int_0^\pi \sin(z \sin \theta - \nu\theta) d\theta$$

$$- \frac{1}{\pi} \int_0^\infty \{ e^{\nu t} + e^{-\nu t} \cos(\nu\pi) \} e^{-z \sinh t} dt \quad (|\arg z| < \frac{1}{2}\pi)$$

9.1.23

$$J_0(x) = \frac{2}{\pi} \int_0^\infty \sin(x \cosh t) dt \quad (x > 0)$$

$$Y_0(x) = -\frac{2}{\pi} \int_0^\infty \cos(x \cosh t) dt \quad (x > 0)$$

9.1.24

$$J_\nu(x) = \frac{2(\frac{1}{2}x)^{-\nu}}{\pi^4 \Gamma(\frac{1}{2}-\nu)} \int_1^\infty \frac{\sin(xt) dt}{(t^2-1)^{\nu+\frac{1}{2}}} \quad (|\Re \nu| < \frac{1}{2}, x > 0)$$

$$Y_\nu(x) = -\frac{2(\frac{1}{2}x)^{-\nu}}{\pi^4 \Gamma(\frac{1}{2}-\nu)} \int_1^\infty \frac{\cos(xt) dt}{(t^2-1)^{\nu+\frac{1}{2}}} \quad (|\Re \nu| < \frac{1}{2}, x > 0)$$

9.1.25

$$H_\nu^{(1)}(z) = \frac{1}{\pi i} \int_{-\infty}^{\infty} e^{z \sinh t - \nu t} dt \quad (|\arg z| < \frac{1}{2}\pi)$$

$$H_\nu^{(2)}(z) = -\frac{1}{\pi i} \int_{-\infty}^{\infty} e^{z \sinh t - \nu t} dt \quad (|\arg z| < \frac{1}{2}\pi)$$

9.1.26

$$J_\nu(x) = \frac{1}{2\pi i} \int_{-t_\infty}^{t_\infty} \frac{\Gamma(-t)(\frac{1}{2}x)^{\nu+2t}}{\Gamma(\nu+t+1)} dt \quad (\Re \nu > 0, x > 0)$$

In the last integral the path of integration must lie to the left of the points  $t=0, 1, 2, \dots$

## Recurrence Relations

$$\begin{aligned} \mathcal{C}_{r-1}(z) + \mathcal{C}_{r+1}(z) &= \frac{2r}{z} \mathcal{C}_r(z) \\ \mathcal{C}_{r-1}(z) - \mathcal{C}_{r+1}(z) &= 2\mathcal{C}'_r(z) \\ \mathcal{C}'_r(z) &= \mathcal{C}_{r-1}(z) - \frac{r}{z} \mathcal{C}_r(z) \\ \mathcal{C}'_r(z) &= -\mathcal{C}_{r+1}(z) + \frac{r}{z} \mathcal{C}_r(z) \end{aligned}$$

$\mathcal{C}$  denotes  $J, Y, H^{(1)}, H^{(2)}$  or any linear combination of these functions, the coefficients in which are independent of  $z$  and  $\nu$ .

$$9.1.27 \quad J'_0(z) = -J_1(z) \quad Y'_0(z) = -Y_1(z)$$

If  $f_r(z) = z^p \mathcal{C}_r(\lambda z^q)$  where  $p, q, \lambda$  are independent of  $\nu$ , then

$$\begin{aligned} f_{r-1}(z) + f_{r+1}(z) &= (2\nu/\lambda) z^{-q} f_r(z) \\ (p+q)f_{r-1}(z) + (p-q)f_{r+1}(z) &= (2\nu/\lambda) z^{1-q} f'_r(z) \\ zf'_r(z) &= \lambda q z^q f_{r-1}(z) + (p+q)f_r(z) \\ zf'_r(z) &= -\lambda q z^q f_{r+1}(z) + (p+q)f_r(z) \end{aligned}$$

## Formulas for Derivatives

$$\left( \frac{1}{z} \frac{d}{dz} \right)^k \{ z^r \mathcal{C}_r(z) \} = z^{r-k} \mathcal{C}_{r-k}(z)$$

$$\left( \frac{1}{z} \frac{d}{dz} \right)^k \{ z^{-r} \mathcal{C}_r(z) \} = (-)^k z^{-r-k} \mathcal{C}_{r+k}(z) \quad (k=0, 1, 2, \dots)$$

$$\begin{aligned} \mathcal{C}^{(k)}(z) &= \frac{1}{2^k} \{ \mathcal{C}_{r-k}(z) - \binom{k}{1} \mathcal{C}_{r-k+2}(z) \\ &\quad + \binom{k}{2} \mathcal{C}_{r-k+4}(z) - \dots + (-)^k \mathcal{C}_{r+k}(z) \} \quad (k=0, 1, 2, \dots) \end{aligned}$$

## Recurrence Relations for Cross-Products

If

$$\begin{aligned} p_r &= J_\nu(a) Y_\nu(b) - J_\nu(b) Y_\nu(a) \\ q_r &= J'_\nu(a) Y'_\nu(b) - J'_\nu(b) Y'_\nu(a) \\ r_r &= J'_\nu(a) Y_\nu(b) - J_\nu(b) Y'_\nu(a) \\ s_r &= J'_\nu(a) Y'_\nu(b) - J'_\nu(b) Y'_\nu(a) \end{aligned}$$

then

$$\begin{aligned} p_{r+1} - p_{r-1} &= -\frac{2r}{a} q_r - \frac{2r}{b} r_r \\ q_{r+1} + r_r &= \frac{v}{a} p_r - \frac{v+1}{b} p_{r+1} \\ r_{r+1} + q_r &= \frac{v}{b} p_r - \frac{v+1}{a} p_{r+1} \\ s_r &= \frac{1}{2} p_{r+1} + \frac{1}{2} p_{r-1} - \frac{v^2}{ab} p_r \end{aligned}$$

and

$$9.1.34 \quad p_r s_r - q_r r_r = \frac{4}{\pi^2 ab}$$

## Analytic Continuation

In 9.1.35 to 9.1.38,  $m$  is an integer.

$$9.1.35 \quad J_\nu(z e^{m\pi i}) = e^{-m\pi i} J_\nu(z) + 2i \sin(m\nu\pi) \cot(\nu\pi) J_\nu'(z)$$

9.1.36

$$Y_\nu(z e^{m\pi i}) = e^{-m\pi i} Y_\nu(z) + 2i \sin(m\nu\pi) \cot(\nu\pi) J_\nu'(z)$$

9.1.37

$$\begin{aligned} \sin(\nu\pi) H_\nu^{(1)}(z e^{m\pi i}) &= -\sin((m-1)\nu\pi) H_\nu^{(1)}(z) \\ &\quad - e^{-m\pi i} \sin(m\nu\pi) H_\nu^{(2)}(z) \end{aligned}$$

9.1.38

$$\begin{aligned} \sin(\nu\pi) H_\nu^{(2)}(z e^{m\pi i}) &= \sin((m+1)\nu\pi) H_\nu^{(2)}(z) \\ &\quad + e^{m\pi i} \sin(m\nu\pi) H_\nu^{(1)}(z) \end{aligned}$$

9.1.39

$$\begin{aligned} H_\nu^{(1)}(z e^{m\pi i}) &= -e^{-m\pi i} H_\nu^{(3)}(z) \\ H_\nu^{(2)}(z e^{-m\pi i}) &= -e^{m\pi i} H_\nu^{(1)}(z) \end{aligned}$$

9.1.40

$$\begin{aligned} J_\nu(\bar{z}) &= \overline{J_\nu(z)} \quad Y_\nu(\bar{z}) = \overline{Y_\nu(z)} \\ H_\nu^{(1)}(\bar{z}) &= \overline{H_\nu^{(2)}(z)} \quad H_\nu^{(2)}(\bar{z}) = \overline{H_\nu^{(1)}(z)} \quad (\nu \text{ real}) \end{aligned}$$

## Generating Function and Associated Series

$$9.1.41 \quad e^{iz(t-1/t)} = \sum_{k=-\infty}^{\infty} t^k J_k(z) \quad (t \neq 0)$$

$$9.1.42 \quad \cos(z \sin \theta) = J_0(z) + 2 \sum_{k=1}^{\infty} J_{2k}(z) \cos(2k\theta)$$

9.1.43

$$\sin(z \sin \theta) = 2 \sum_{k=0}^{\infty} J_{2k+1}(z) \sin((2k+1)\theta)$$

9.1.44

$$\cos(z \cos \theta) = J_0(z) + 2 \sum_{k=1}^{\infty} (-)^k J_{2k}(z) \cos(2k\theta)$$

9.1.45

$$\sin(z \cos \theta) = 2 \sum_{k=0}^{\infty} (-)^k J_{2k+1}(z) \cos((2k+1)\theta)$$

9.1.46

$$1 = J_0(z) + 2J_2(z) + 2J_4(z) + 2J_6(z) + \dots$$

9.1.47

$$\cos z = J_0(z) - 2J_2(z) + 2J_4(z) - 2J_6(z) + \dots$$

$$9.1.48 \quad \sin z = 2J_1(z) - 2J_3(z) + 2J_5(z) - \dots$$



9.1.72

$$\lim \{v^\nu Q_{\nu}^{-\mu} \left( \cos \frac{x}{v} \right)\} = -\frac{1}{2}\pi Y_\nu(x) \quad (x>0)$$

For  $P_\nu^{-\mu}$  and  $Q_\nu^{-\mu}$ , see chapter 8.

## Continued Fractions

9.1.73

$$\begin{aligned} J_\nu(z) &= \frac{1}{2\nu z^{-1}} - \frac{1}{2(\nu+1)z^{-1}} - \frac{1}{2(\nu+2)z^{-1}} - \dots \\ &= \frac{\frac{1}{2}z/\nu}{1-\frac{1}{1-\frac{\frac{1}{2}z^2/\{\nu(\nu+1)\}}{1-\frac{1}{1-\frac{\frac{1}{2}z^2/(\nu+1)(\nu+2)}{\dots}}}}} \end{aligned}$$

## Multiplication Theorem

9.1.74

$$\mathcal{C}_r(\lambda z) = \lambda^{\pm\nu} \sum_{k=0}^{\infty} \frac{(\mp)^k (\lambda^2 - 1)^k (\frac{1}{2}z)^k}{k!} \mathcal{C}_{r\pm k}(z) \quad (|\lambda^2 - 1| < 1)$$

If  $\mathcal{C}=J$  and the upper signs are taken, the restriction on  $\lambda$  is unnecessary.This theorem will furnish expansions of  $\mathcal{C}_r(re^{i\theta})$  in terms of  $\mathcal{C}_{r\pm k}(r)$ .

## Addition Theorems

Neumann's

$$9.1.75 \quad \mathcal{C}_r(u \pm v) = \sum_{k=-\infty}^{\infty} \mathcal{C}_{r\mp k}(u) J_k(v) \quad (|v| < |u|)$$

The restriction  $|v| < |u|$  is unnecessary when  $\mathcal{C}=J$  and  $v$  is an integer or zero. Special cases are

$$9.1.76 \quad 1 = J_0^2(z) + 2 \sum_{k=1}^{\infty} J_k^2(z)$$

9.1.77

$$0 = \sum_{k=0}^{2n} (-1)^k J_k(z) J_{2n-k}(z) + 2 \sum_{k=1}^{\infty} J_k(z) J_{2n+k}(z) \quad (n \geq 1)$$

9.1.78

$$J_n(2z) = \sum_{k=0}^n J_k(z) J_{n-k}(z) + 2 \sum_{k=1}^{\infty} (-1)^k J_k(z) J_{n+k}(z)$$

Grafs

9.1.79

$$\mathcal{C}_r(w) \frac{\cos \nu x}{\sin} = \sum_{k=-\infty}^{\infty} \mathcal{C}_{r+k}(u) J_k(v) \frac{\cos k\alpha (|ve^{\pm i\alpha}| < |u|)}{\sin}$$


Gegenbauer's

9.1.80

$$\frac{\mathcal{C}_r(w)}{w^\nu} = 2^\nu \Gamma(\nu) \sum_{k=0}^{\infty} (\nu+k) \frac{\mathcal{C}_{r+k}(u)}{u^\nu} \frac{J_{r+k}(v)}{v^\nu} C_k^{(\nu)}(\cos \alpha) \quad (\nu \neq 0, -1, \dots, |ve^{\pm i\alpha}| < |u|)$$

In 9.1.79 and 9.1.80,  
 $w = \sqrt{(u^2 + v^2 - 2uv \cos \alpha)}$ ,

$$u - v \cos \alpha = w \cos \chi, \quad v \sin \alpha = w \sin \chi$$

the branches being chosen so that  $w \rightarrow u$  and  $x \rightarrow 0$  as  $v \rightarrow 0$ .  $C_k^{(\nu)}(\cos \alpha)$  is Gegenbauer's polynomial (see chapter 22).

Gegenbauer's addition theorem.

If  $u, v$  are real and positive and  $0 \leq \alpha \leq \pi$ , then  $w, x$  are real and non-negative, and the geometrical relationship of the variables is shown in the diagram.The restrictions  $|ve^{\pm i\alpha}| < |u|$  are unnecessary in 9.1.79 when  $\mathcal{C}=J$  and  $\nu$  is an integer or zero, and in 9.1.80 when  $\mathcal{C}=J$ .Degenerate Form ( $u = \infty$ ):

9.1.81

$$e^{iz \cos \alpha} = \Gamma(\nu) (\frac{1}{2}v)^{-\nu} \sum_{k=0}^{\infty} (\nu+k) i^k J_{r+k}(v) C_k^{(\nu)}(\cos \alpha) \quad (\nu \neq 0, -1, \dots)$$

Neumann's Expansion of an Arbitrary Function in a Series of Bessel Functions

$$9.1.82 \quad f(z) = a_0 J_0(z) + 2 \sum_{k=1}^{\infty} a_k J_k(z) \quad (|z| < c)$$

where  $c$  is the distance of the nearest singularity of  $f(z)$  from  $z=0$ ,

$$9.1.83 \quad a_k = \frac{1}{2\pi i} \int_{|z|=c'} f(t) O_k(t) dt \quad (0 < c' < c)$$

and  $O_k(t)$  is Neumann's polynomial. The latter is defined by the generating function

9.1.84

$$\frac{1}{t-z} = J_0(z) O_0(t) + 2 \sum_{k=1}^{\infty} J_k(z) O_k(t) \quad (|z| < |t|)$$

 $O_n(t)$  is a polynomial of degree  $n+1$  in  $1/t$ ;  $O_0(t) = 1/t$ ,

9.1.85

$$O_n(t) = \frac{1}{4} \sum_{k=0}^{\leq n} \frac{n(n-k-1)!}{k!} \left(\frac{2}{t}\right)^{n-2k+1} \quad (n=1, 2, \dots)$$

The more general form of expansion

$$9.1.86 \quad f(z) = a_0 J_0(z) + 2 \sum_{k=1}^{\infty} a_k J_{r+k}(z)$$

	$f(s)$		$F(t)$
29.3.126	$e^{as} E_1(as)$	$(a > 0)$	5 $\frac{1}{t+a}$
29.3.127	$\frac{1}{a} - se^{as} E_1(as)$	$(a > 0)$	5 $\frac{1}{(t+a)^2}$
29.3.128	$a^{1-n} e^{at} E_n(as)$	$(a > 0; n=0, 1, 2, \dots)$	5 $\frac{1}{(t+a)^n}$
29.3.129	$\left[ \frac{\pi}{2} - \text{Si}(s) \right]$	$\cos s + \text{Ci}(s) \sin s$	5 $\frac{1}{t^2+1}$

29.4. Table of Laplace-Stieltjes Transforms<sup>4</sup>

	$\Phi(s)$		$\Phi(t)$
29.4.1	$\int_0^{\infty} e^{-st} d\Phi(t)$		$\Phi(t)$
29.4.2	$e^{-kt}$	$(k > 0)$	$u(t-k)$
29.4.3	$\frac{1}{1-e^{-ks}}$	$(k > 0)$	$\sum_{n=0}^{\infty} u(t-nk)$
29.4.4	$\frac{1}{1+e^{-ks}}$	$(k > 0)$	$\sum_{n=0}^{\infty} (-1)^n u(t-nk)$
29.4.5	$\frac{1}{\sinh ks}$	$(k > 0)$	$2 \sum_{n=0}^{\infty} u[t-(2n+1)k]$
29.4.6	$\frac{1}{\cosh ks}$	$(k > 0)$	$2 \sum_{n=0}^{\infty} (-1)^n u[t-(2n+1)k]$
29.4.7	$\tanh ks$	$(k > 0)$	$u(t) + 2 \sum_{n=1}^{\infty} (-1)^n u(t-2nk)$
29.4.8	$\frac{1}{\sinh (ks+a)}$	$(k > 0)$	$2 \sum_{n=0}^{\infty} e^{-(2n+1)a} u[t-(2n+1)k]$
29.4.9	$\frac{e^{-ht}}{\sinh (ks+a)}$	$(k > 0, h > 0)$	$2 \sum_{n=0}^{\infty} e^{-(2n+1)a} u[t-h-(2n+1)k]$
29.4.10	$\frac{\sinh (hs+b)}{\sinh (ks+a)}$	$(0 < h < k)$	$\sum_{n=0}^{\infty} e^{-(2n+1)a} \{ e^b u[t+h-(2n+1)k] - e^{-b} u[t-h-(2n+1)k] \}$
29.4.11	$\sum_{n=0}^{\infty} a_n e^{-kn^2}$	$(0 < k_0 < k_1 < \dots)$	$\sum_{n=0}^{\infty} a_n u(t-k_n)$

For the definition of the Laplace-Stieltjes transform see [29.7]. In practice, Laplace-Stieltjes transforms are often written as ordinary Laplace transforms involving Dirac's delta function  $\delta(t)$ . This "function" may formally be considered as

the derivative of the unit step function,  $du(t) = \delta(t)$   $dt$ , so that  $\int_{-\infty}^x du(t) = \int_{-\infty}^x \delta(t) dt = \begin{cases} 0 & (x < 0) \\ 1 & (x > 0) \end{cases}$ . The correspondence 29.4.2, for instance, then assumes the form  $e^{-kt} = \int_0^{\infty} e^{-st} \delta(t-k) dt$ .

<sup>4</sup> Adapted by permission from P. M. Morse and H. Feshbach, Methods of theoretical physics, vols. 1, 2, McGraw-Hill Book Co., Inc., New York, N.Y., 1953.

# 9. Bessel Functions of Integer Order

## Mathematical Properties

is chapter are for Bessel functions of integer order; the text treats generalizations used are:

1. integer or zero.  
2. except where otherwise stated real in the sections devoted to **9.9, 9.10, and 9.11.**

3. for the Bessel functions is [15] and the British Association Mathematical Tables. The function denoted  $N_v(z)$  by physicists is

are those of:

### Bessel Functions $J$ and $Y$

#### 9.1. Definitions and Elementary Properties

##### Differential Equation

$$9.1.1 \quad z^2 \frac{d^2w}{dz^2} + z \frac{dw}{dz} + (z^2 - \nu^2)w = 0$$

Solutions are the Bessel functions of the first kind  $J_{\pm\nu}(z)$ , of the second kind  $Y_\nu(z)$  (also called Weber's function) and of the third kind  $H_\nu^{(1)}(z), H_\nu^{(2)}(z)$  (also called the Hankel functions). Each is a regular (holomorphic) function of  $z$  throughout the  $z$ -plane cut along the negative real axis, and for fixed  $z (\neq 0)$  each is an entire (integral) function of  $\nu$ . When  $\nu = \pm n$ ,  $J_\nu(z)$  has no branch point and is an entire (integral) function of  $z$ .

# Dynamic dictionary of math functions

## Computer algebra:

- **classic**: polynomial to represent their roots + following tools: euclidian division, Euclid algorithm, Gröbner bases.
- **modern**: linear differential equation as data structures to represent their solutions [SaZi94, ChSa98, Chyzak00, MeSa03, Salvy05] with same tools as classical case but non-commutative.
- **prototype** ESF at <http://algo.inria.fr/esf> (65% of Abramowitz-Stegun)
- **todo**: interactivity, integral transforms, parametric integrals.



# ReActivity

Wendy Mackay, INRIA Saclay,  
J.-D. Fekete, INRIA Saclay,  
Mary Czerwinski, MSRR,  
George Robertson, MSRR

Michel Beaudouin-Lafon, Paris 11,  
Olivier Chapuis, CNRS,  
Pierre Dragicevic, INRIA Saclay,  
Emmanuel Pietriga, INRIA Saclay,  
Aurélien Tabard, Paris 11 (PhD)

## Logs of experiments for biologists, historians, other scientists

- mixed inputs from lab notebooks and computers,
- interactive visualization of scientific activity,
- support for managing scientific workflow.



CENTRE DE RECHERCHE  
COMMUN

INRIA  
MICROSOFT RESEARCH

# ReActivity

## Recent results:

- workshop on Interacting with **Temporal Data** at CHI'09 (35 participants)
- streamlining the computation of aggregated metrics on Wikipedia “live” and small focused “Dashboard visualizations” tools
- WILD: Wall-sized **Interaction** with Large Datasets (32 screens, 8 Vicon, 1 interactive table expansions)
- Augmented Paper/Electronic Notebooks

# Adaptive Combinatorial Search for E-science

Youssef Hamadi, MSRC

Marc Schoenauer, INRIA-Saclay

Anne Auger, INRIA-Saclay

Lucas Bordeaux, MSRC


Michèle Sebag, CNRS

## Parallel constraint programming and optimization for very large scientific data

- improve **the usability** of *Combinatorial Search* algorithms.
- automate the fine tuning of solver parameters.
- parallel solver: “disolver”



MoGo



CENTRE DE RECHERCHE  
COMMUN

INRIA  
MICROSOFT RESEARCH

# Adaptive Combinatorial Search for E-science

- **constraint programming**: learn instance-dependent variable ordering
- **evolutionary algorithms**: use multi-armed bandit algorithms and extreme values statistics
- **continuous search spaces**: use local curvature

# Image and video mining for science and humanities

Jean Ponce, ENS


Andrew Blake, MSRC

Patrick Pérez, INRIA Rennes

Cordelia Schmid, INRIA Grenoble

## Computer vision and Machine learning for:

- *sociology*: human activity modeling and recognition in video archives
- *archaeology and cultural heritage preservation*: 3D object modeling and recognition from historical paintings and photographs
- environmental sciences: change detection in dynamic satellite imagery




CENTRE DE RECHERCHE  
COMMUN

INRIA MICROSOFT RESEARCH

# Sciences in track B

DDMF	computer algebra	hard sciences
Adapt. search	constraints, machine learning	hard sciences, biology
Reactivity	chi + visualisation	soft sciences, biology
I.V. mining	computer vision	humanities, environment



# Future

CENTRE DE RECHERCHE  
COMMUN



# Objectives

- 30 resident researchers
- tight links with French academia (phD, post-doc)
- develop useful research for scientific community
- provide public tools (BSD-like license)
- become a new and attractive pole in CS research
- and source of spin off companies



- vision and medical applications ?



CENTRE DE RECHERCHE  
COMMUN



INRIA  
MICROSOFT RESEARCH